

The background of the page is a 3D molecular model. It features several large, semi-transparent spheres in shades of red, blue, and white, connected by thin, metallic-looking rods. The spheres are arranged in a complex, branching structure, resembling a protein chain or a network of atoms. The background is a light, hazy blue with scattered, smaller, out-of-focus spheres in various colors, creating a sense of depth and scientific atmosphere.

Dietary Protein and the Critical Care Patient



Dr Keith Garleb, PhD, Chief Science Officer, Garleb Brothers Consulting, and Adjunct Professor University of Illinois, USA

Proteins are macromolecules composed of dispensable and indispensable amino acids. The quality of a protein is a function of its digestibility and indispensable amino acid content. Critical care patients have an increased need for protein to meet their daily nitrogen and amino acid requirements. Delivering high quality protein to this population is crucial for their recovery.

What is protein?

Proteins are large molecules made up of amino acids linked together by peptide bonds. There are 20 amino acids.¹ Amino acids can be divided into two groups: indispensable (essential) and dispensable (non-essential). Indispensable amino acids cannot be synthesized by the body and must be obtained through the diet. There are nine indispensable amino acids, which include: histidine, isoleucine, leucine, lysine, methionine, phenylalanine, threonine, tryptophan and valine. Dietary requirements

for indispensable amino acids are presented in **Table 1**. The remaining amino acids (alanine, aspartic acid, asparagine, glutamic acid, serine, arginine, glutamine, glycine, proline, cysteine and tyrosine) are considered dispensable because they can be synthesized by the body. However, under specific physiological or pathological conditions some of the dispensable amino acids can become conditionally indispensable because endogenous synthesis cannot meet metabolic needs.² These include: arginine, cysteine, glutamine, glycine, proline and tyrosine.

“Similar to a chain’s strength being limited by its weakest link, a protein’s capacity to meet nutritional needs is limited by its ‘most deficient’ indispensable amino acid.”

Table 1: Recommended Daily Allowance of Indispensable Amino Acids for Adults 19 years and Older¹

Amino acid	mg/kg/day
Histidine	14
Isoleucine	19
Leucine	42
Lysine	38
Methionine (+ cysteine)	19
Phenylalanine (+ tyrosine)	33
Threonine	20
Tryptophan	5
Valine	24

Once ingested, the low pH of the stomach denatures the protein which is then broken down into peptides by the action of pepsin. After moving into the small intestine, further digestion takes place through the action of pancreatic enzymes, such as trypsin, chymotrypsin and carboxypeptidase, forming free amino acids, dipeptides and tripeptides that can be absorbed. The digestibility of protein can vary. Animal proteins tend to have a higher digestibility (~95%) compared to whole plant foods (~80-85%).³ However, purified plant protein sources such as soy protein isolate and pea protein concentrate display a digestibility similar to that of animal-based protein sources.⁴ Protein sources can also vary significantly in their content of amino acids. A protein is considered complete if it contains an adequate proportion of each of the nine indispensable amino acids (e.g. milk protein). Plant proteins are generally lower in methionine (e.g. legumes) and lysine (e.g. cereals) than animal-based proteins.^{5, 6} The indispensable amino acid that is most deficient relative to metabolic needs is referred to as the limiting amino acid within the protein. Similar to a chain’s strength being limited by its weakest link, a protein’s capacity to meet nutritional needs is limited by its ‘most deficient’ indispensable amino acid. To overcome the consequence of a limiting amino acid, the amino acid can be fortified as is the case with methionine in soy-based infant formulas. In addition, proteins can be combined to complement their amino acid profiles and provide a more complete protein blend.^{5, 6}

Protein quality

Protein digestibility and content of indispensable amino acids are factors that affect protein quality. Over the years, numerous methods have been developed in an attempt to quantify the quality of a

given protein. One of the methods, the Protein Digestibility-Corrected Amino Acid Score (PDCAAS), is now widely used as a routine assay for the assessment of the quality of single proteins and protein blends.^{7, 8} According to this method, protein quality is assessed by expressing the amount of the first limiting indispensable amino acid in the protein of interest as a fraction of the same amino acid in an ‘ideal’ reference pattern and then multiplying this amino acid score by the true faecal nitrogen digestibility (%) as determined in rats. Scores exceeding 100% are truncated to 100%. Higher scores reflect a higher quality protein or protein blend.

Protein requirements

The requirement for dietary protein is based on the body’s need for nitrogen and indispensable amino acids. Nitrogen is needed for the synthesis of dispensable and conditionally indispensable amino acids, as well as physiologically important nitrogen containing compounds such as nucleic acids. As previously stated, indispensable amino acids cannot be made by human tissues at a rate commensurate with metabolic needs and must be supplied via the diet. Unlike carbohydrate (glycogen) and fat (adipose) stores, whose primary role is energy, the protein lost during fasting or periods of physical stress has a greater negative impact on the body due to the specific and relevant roles that protein plays. During periods of high physical stress, such as illness or hospitalisation, patients become hypercatabolic and may experience a significant loss of muscle protein as a result of anabolic resistance and an increase in muscle protein breakdown.⁹ Inadequate protein intake can lead to a negative nitrogen balance resulting in skeletal muscle atrophy, functional decline and poor recoveries.¹⁰

The administration of high quality proteins (high PDCAAS score) can help mitigate the metabolic loss of protein during periods of physical stress by providing the nitrogen and indispensable amino acids needed by the body for recovery.

According to the Institute of Medicine (2005), adults require 0.8 grams protein/kg body weight/day with requirements being higher for pregnant and lactating women and growing children.¹ There is considerable interest and effort to better understand the protein/amino acid requirements of hospitalised patients, particularly those who are critically ill. Following a review of the literature, researchers concluded that critically ill patients require significant amounts of protein, likely in excess of 1.2 grams/kg body weight/day to maintain protein balance and lean body mass.¹¹ Current recommendations suggest that 1.2-1.5 grams protein/kg body weight/day is sufficient for most critically ill patients.^{12, 13, 14} The European Society for Enteral and Parenteral Nutrition (ESPEN) recommends from 1.3-1.5 grams protein/kg body weight/day for most critical care patients.¹³ The Society of Critical Care Medicine (SCCM) and the American Society of Parenteral and Enteral Nutrition (ASPEN) recommend a protein intake of 1.2-2.0 grams/kg body weight/day. Older

patients in intensive care units appear less responsive to lower protein intakes and may require more than 2 grams/kg body weight/day.^{15, 16} A systematic review by Hoffer and Bistrian (2012) suggests that 2.0-2.5 grams protein/kg body weight/day is safe and could be optimum for most critically ill patients.¹⁷ While protein recommendations vary for critically ill patients, all are significantly greater than the 0.8 grams/kg body weight/day recommended for healthy adults. Unfortunately, results from an international multicenter observational study show that critically ill adults receive only about 0.6 grams protein/kg body weight/day which is well below the levels recommended by authoritative bodies.¹⁸

Conclusion

Adequate protein is essential for the recovery of critically ill patients. Due to their condition, protein requirements are substantially higher than those of healthy individuals. In order to meet their demands, increased intakes of high-quality protein blends that can be easily delivered to the patient are warranted. High quality proteins, as indicated by their PDCAAS score, ensure the delivery of a highly digestible protein blend that provides the nitrogen and indispensable amino acids essential for recovery.

“High quality proteins, as indicated by their PDCAAS score, ensure the delivery of a highly digestible protein blend that provides the nitrogen and indispensable amino acids essential for recovery.”

About the author: Dr Garleb is dedicated to the discovery and creation of nutrition technologies that impact health and wellness and improve patient outcomes. Dr. Garleb has 20 patents and over 100 peer reviewed articles, abstracts and reviews to his credit.

Dr Garleb has a Bachelor of Science in Agricultural Sciences, a Master of Science in Animal Sciences and a PhD in Nutritional Biochemistry from the University of Illinois at Urbana-Champaign. He is currently an Adjunct Professor at the University.

References: **1.** Institute of Medicine (2005). Dietary reference intakes for energy, carbohydrate, fiber, fat, fatty acids, cholesterol, protein, and amino acids. Washington, DC: The National Academies Press. **2.** Laidlaw SA, Kopple JD (1987). Newer concepts of the indispensable amino acids. *Am J Clin Nutr*; 46(4): 593-605. **3.** Wu G (2016). Dietary protein intake and human health. *Food Funct*; 7(3): 1251-1265. **4.** FAO (2012). Report of a sub-committee of the 2011 FAO consultation on 'Protein Quality Evaluation in Human Nutrition': the assessment of amino acid digestibility in foods for humans and including a collation of published ileal amino acid digestibility data from human foods. Rome (Italy). **5.** van Vliet S, et al. (2015). The skeletal muscle anabolic response to plant- versus animal-based protein consumption. *J Nutr*; 145(9): 1981-1991. **6.** Gorissen SHM, et al. (2018). Protein content and amino acid composition of commercially available plant-based protein isolates. *Amino Acids*; 50(12):1685-1695. **7.** FAO/WHO Expert Consultation (1991). Protein quality evaluation. Food and Agriculture organization of the United Nations. FAO Food and Nutrition Paper. No. 51, Rome. **8.** Schaafsma G (2012). Advantages and limitations of the protein digestibility-corrected amino acid score (PDCAAS) as a method for evaluating protein quality in human diets. *Br J Nutr*; 108(2): S333-336. **9.** Berg A, et al. (2013). Whole body protein kinetics during hypocaloric and normocaloric feeding in critically ill patients. *Critical Care*; 17(4): R158. **10.** Wejls PJM, et al. (2014). Protein and amino acids are fundamental to optimal nutrition support in critically ill patients. *Critical Care*; 18: 591. **11.** Wejls PJM, Wischmeyer PE (2013). Optimizing energy and protein balance in the ICU. *Curr Opin Clin Nutr Metab Care*; 16(2): 194-201. **12.** Ziegler TR (2009). Parenteral nutrition in the critically ill. *N Engl J Med*; 361(11): 1088-1097. **13.** Singer P, et al. (2009). ESPEN guidelines on parenteral nutrition: intensive care. *Clin Nutr*; 28(4): 387-400. **14.** McClave SA, et al. (2009). Guidelines for the provision and assessment of nutrition support therapy in the adult critically ill patient: Society of Critical Care Medicine (SCCM) and American Society for Parenteral and Enteral Nutrition (ASPEN). *JPEN*; 33(3): 277-316. **15.** Dickerson RN, et al. (2015). Influence of aging on nitrogen accretion during critical illness. *JPEN*; 39(3): 282-290. **16.** Dickerson RN (2016). Nitrogen balance and protein requirements for critically ill older patients. *Nutrients*; 8(4): 226. **17.** Hoffer LJ, Bistrian BR (2012). Appropriate protein provision in critical care illness: A systemic and narrative review. *Am J Clin Nutr*; 96(3): 591-600. **18.** Alberda C, et al. (2009). The relationship between nutritional intake and clinical outcomes in critically ill patients: results of an international multicenter observational study. *Intensive Care Med*; 35(10): 1728-1737.



NOW TEST YOUR KNOWLEDGE
Visit CPD section at: www.nutrition2me.com

The CNPD questionnaire linked to this article has been kindly sponsored by **nutrinovo**: www.nutrinovo.com

nutrinovo
NUTRITION TECHNOLOGIES



WHY COMPROMISE?

**COMPLETE
IN ESSENTIAL
AMINO ACIDS**

**11G
PROTEIN
IN 45ML**

**ISOTONIC
OSMOLALITY
LEVEL**

**HIGHEST
AMINO ACID
CONCENTRATION**

ProSource TF represents a unique and unrivalled option in liquid proteins designed specifically to support enteral feeding. Not only is TF complete in ALL essential amino acids, but it contains those amino acids at a far higher concentration than other liquid proteins.

ProSource TF PDCAAS Score = 100

Essential Amino Acids	IOM Reference Pattern (mg/g protein)	ProSource TF Pattern (mg/g protein)
Histidine	17	18
Isoleucine	23	24
Leucine	52	55
Lysine	47	51
Methionine & Cysteine	23	24
Phenylalanine & Tyrosine	41	43
Threonine	26	28
Tryptophan	6	6
Valine	29	31

Each gram of TF protein delivers more of what your patients need, without compromising on volume, tolerance and quality.



For the Dietary
Management of
Hypoproteinaemia.

PROSource

A High Protein Liquid Supplement **TF**

11 grams of protein
per 45 ml

Try ProSource TF for free www.nutrinovo.com

nutrinovo
simply innovative nutrition