

Request a **FREE sample pack of any ProSource product – direct to you or direct to your patient.**

- Sample packs can be delivered direct to you or direct to your patient in the community.
- The majority of sample packs are despatched within 24 hours and we aim to despatch 100% within 48 hours, meaning you or your patients are able to try any ProSource product when you need it most.

**FREE
SAMPLE
SERVICE**

Request a **FREE** sample pack, visit www.nutrinovo.com
email info@nutrinovo.com or call **01249 691301**



nutrinovo
simply innovative nutrition

Nutrino Limited | +44 (0)1249 691301

Manufactured in the USA for Nutrinovo Limited.

www.nutrinovo.com @Nutrinovo

WHY COMPROMISE?



ProSource TF represents a unique and unrivalled option in liquid protein designed specifically to support enteral feeding. Not only is TF complete in ALL essential amino acids, but it contains those amino acids at a far higher concentration than other liquid proteins.

COMPLETE IN ESSENTIAL AMINO ACIDS

ProSource TF contains all essential amino acids, at levels that closely match the US Institute of Medicine's profile for an "ideal protein".

HIGHEST AMINO ACID CONCENTRATION

ProSource TF contains the essential amino acids at levels of around double that of other manufacturer's liquid protein supplements. This means that each gram of TF protein delivers more of what your patients need, without compromise.

ISOTONIC OSMOLALITY LEVEL

ProSource TF has an osmolality level of 380 mOsm/kg water, providing a close to isotonic osmolality level, thus helping to improve patient tolerance.

11G PROTEIN IN 45ML

Enables you to increase the protein intake of patients being enterally tube fed, without significantly increasing their carbohydrates or volume; or changing their required enteral feed.

ACBS APPROVED

ProSource TF is ACBS approved for Hypoproteinaemia. Hypoproteinaemia occurs as a result of many disease states, meaning that ProSource TF can be useful across a wide range of patient conditions.



EASE OF ADMINISTRATION

The presentation of ProSource TF allows it to be administered in a way that best suits each patient's need - either directly through an enteral feeding tube or via an enteral giving set.

Below is a suggested method of administration which is quick, easy, limits any wastage and reduces contamination risk.

1. Remove the plunger from a 60ml enteral syringe
2. Connect the enteral syringe to the patients enteral feeding tube or giving set
3. Open the sachet of TF by removing the easy to tear tab from the top
4. Pour the contents directly into the top of the syringe

We recommend flushing the patient's feeding tube with water both before and after the addition of ProSource TF.

